

## Errata and Updates for ASM Exam MAS-I (First Edition Second Printing) Sorted by Page

Practice Exam 5 Question 6 is defective. See the correction to the question below. In Practice Exam 6 Question 11, the answer ranges are suitable for an annuity of 1 per year rather than 50,000 per year.

[8/31/2018] On page 153, in the solution to exercise 13.30, on the first displayed line, change  $10 + 7e^{-(10-x)/7}$  to  $10 - 7e^{-(10-x)/7}$ .

[10/28/2018] On page 159, in the solution to exercise 14.9, on the first line, change “with mean 1” to “on  $[0, 1]$ ”.

[10/28/2018] On page 206, two lines after formula (18.2), change  $1 - 10(0.9^9) = 0.964844$  to  $1 - 10(0.5^9) = 0.980469$ .

[10/28/2018] On page 292, in the box before exercises 24.8 and 24.9, on the third line, change “geometric” to “gamma”.

[7/20/2018] On page 293, in the solution to exercise 24.1, change the answer key from **(C)** to **(B)**.

[7/1/2018] On page 293, in the solution to exercise 24.3, on the sixth line (the one with  $u \rightarrow$ ), change  $\sqrt{(1-u)/2}$  to  $1 - \sqrt{(1-u)/2}$ . Replace the last two lines of the solution with

Comparing that fraction to the second number in each pair, we have  $0.55 \leq 1 - 0.3 = 0.7$ ,  $0.37 \leq 1 - 0.4 = 0.6$ ,  $0.62 \geq 0.6$ , and  $0.77 \leq 0.8$ . We accept when the second number is less than or equal to the fraction, so we accept all except the third number. The average is  $(0.3 + 0.4 + 0.8)/3 = \boxed{0.5}$ .

[10/28/2018] On page 295, change the first two lines of the solution to exercise 24.7 to

$$c = \max\left(\frac{f(x)}{g(x)}\right)$$

$$\frac{f(x)}{g(x)} = \frac{xe^{-x}}{1/(1+x)^2}$$

[10/28/2018] On page 314, in the solution to exercise 25.24, on the third line, change “by  $\theta$ ” to “be  $\theta$ ”.

[11/6/2018] On page 330, on lines 6 through 9 from the bottom of the page, add a factor  $b$  before each  $k$ :

$$bk_5(16) = \frac{e^{-((16-5)^2)/2(8^2)}}{\sqrt{2\pi}} = 0.155012$$

$$bk_{12}(16) = \frac{e^{-((16-12)^2)/2(8^2)}}{\sqrt{2\pi}} = 0.352065$$

$$bk_{15}(16) = \frac{e^{-((16-15)^2)/2(8^2)}}{\sqrt{2\pi}} = 0.395838$$

$$bk_{20}(16) = \frac{e^{-((16-20)^2)/2(8^2)}}{\sqrt{2\pi}} = 0.352065$$

[7/29/2018] On page 418, in exercise 29.25, on the second line, the square symbol should be on the denominator:

$$S_X(x) = \frac{\theta^4}{(\theta^2 + x^2)^2}$$

[10/28/2018] On page 549, in exercise 36.22, in the third bullet, change all 4 denominators from  $n$  to  $\sqrt{n}$ .

- [10/28/2018] On page 599, in the solution to exercise 40.3, on the fifth line, change  $\lambda^{-n\bar{X}}$  to  $\lambda^{n\bar{X}}$ .
- [10/28/2018] On page 601, in the solution to exercise 40.9, on the second-to-last line, change the  $-2$  at the beginning of the line and the  $-2$  after “or” to  $2$ ; remove the minus signs.
- [10/28/2018] On page 603, on the third line of the third paragraph, change “20<sup>th</sup> percentile” to “15<sup>th</sup> percentile”.
- [10/28/2018] On page 606, in the solution to exercise 41.4, on the first line, change  $(3, 8)$  to  $(5, 8)$ .
- [7/6/2018] On page 635, in the fourth displayed line, on the right side in the exponent, change  $\sum_{i=1}^p$  to  $\sum_{i=2}^p$ .
- [8/12/2018] On page 642, change the first two lines of exercise 44.15 to

In a cumulative proportional odds model for an ordinal variable, the fitted model is

$$\ln \frac{\sum_{i=1}^j \hat{\pi}_i}{1 - \sum_{i=1}^j \hat{\pi}_i} = b_{0j} + b_1 x_1$$

- [8/12/2018] On page 649, in the solution to exercise 44.23, two lines from the end, change  $\Phi(1.01)$  to  $\Phi(1.32)$ .
- [7/10/2018] On page 682, on the third line of Quiz 46-4, change “loglikelihood” to “loglikelihood”.
- [6/28/2018] On page 704, in exercise 47.20, in the first bullet, change the two subscripts on the right to  $i_2$  and  $i_3$ :
- $$\hat{y}_i = 20.0 - 1.5x_{i_2} - 2.0x_{i_3}$$
- [6/28/2018] On page 707, in the solution to exercise 47.12, on the third line, change  $b_2$  to  $b_3$ .
- [6/28/2018] On page 712, on the first line of Example 48B, change  $\beta_5 x_6$  to  $\beta_5 x_5$ .
- [6/28/2018] On page 715, in exercise 48.2, on the second line, replace  $x_{i_7}$  with  $x_{i_6}$ .
- [7/17/2018] On page 719, in exercise 48.15, on the second line, replace  $\beta_4 x_{i_1} x_{i_2}$  with  $\beta_4 x_{i_2} x_{i_3}$ .
- [8/8/2018] On page 719, exercise 48.16 is a duplicate of exercise 48.13.
- [8/8/2018] On page 720, in exercise 48.17, on the second line of the second bullet, put a period after 0.940. The statement beginning “A second regression equation” should be moved to a third bullet.
- [6/28/2018] On page 721, in exercise 48.21, on the last line, change  $\beta_3 = 0$  to  $\beta_4 = 0$ .
- [8/8/2018] On page 727, in the solutions to exercises 48.10 and 48.11, on the displayed line in each solution,  $SSE_R$  and  $SSE_{UR}$  should be interchanged so that the numerator is  $(SSE_R - SSE_{UR})/q$ . In addition, in the solution to exercise 48.11, put parentheses around  $n - p$  in the denominator.
- [8/8/2018] On page 727, in the solution to exercise 48.13, on the first and second lines, delete the sentence fragment “The unrestricted model with 8 variables.”
- [8/8/2018] On page 727, in the solution to exercise 48.14, on the displayed line, change  $n - k$  to  $n - p$ .
- [8/8/2018] On page 737, in exercise 49.2, in the first bullet, change  $\hat{y}$  to  $\hat{y}_i$ .
- [8/8/2018] On page 737, in the solution to exercise 49.2, on the second line, change  $\hat{y}$  to  $\hat{y}_i$ . Replace the last line with

$$r_3 = \frac{\hat{\varepsilon}_3}{s\sqrt{1 - h_{33}}} = \frac{1.2}{2\sqrt{1 - 0.6}} = \boxed{0.948683}$$

- [6/28/2018] On page 743, two lines above equation (50.1), change  $\beta_0 + \beta_1 x^*$  to  $\beta_1 + \beta_2 x^*$ .
- [7/10/2018] On page 791, in the solution to exercise 52.5, on the first line, delete “a” before “Poisson”.

- [7/2/2018] On page 804, one line below formula (54.1), change “variance of the residual  $\varepsilon$ ” to “residual variance of the regression”.
- [7/2/2018] On page 805, in the first paragraph of Example 54A, change “The variance of the residual” to “The residual variance of the regression”.
- [7/2/2018] On page 807, in exercises 54.8 and 54.9, change “estimated variance of the residuals” to “estimated residual variance of the regression”.
- [7/2/2018] On page 808, in exercise 54.10, 2–3 lines from the end, delete the sentence beginning “For each model”.
- [7/2/2018] On page 808, in exercise 54.11, on the first line, change 28 to 29. In the table, delete the  $\hat{\sigma}^2$  column.
- [8/8/2018] On page 811, in the solution to exercise 54.2, on the second line, replace “6 models” with “7 models” and replace “5” with “6”.
- [8/8/2018] On pages 811–812, replace the solution to exercise 54.10 with
- The mean squared error of the full model is  $284/(60 - 5) = 5.163636$ . Then

$$C_p(0) = \frac{326}{60} = 5.4333$$

$$C_p(1) = \frac{314 + 2(5.163636)}{60} = 5.4055$$

$$C_p(2) = \frac{303 + 2(2)(5.163636)}{60} = 5.3942$$

$$C_p(3) = \frac{293 + 2(3)(5.163636)}{60} = 5.4000$$

$$C_p(4) = \frac{284 + 2(4)(5.163636)}{60} = 5.4218$$

The **2-variable** model is selected.

- [8/8/2018] On page 812, replace the solution to exercise 54.11 with
- The estimated value of the mean square error of the model with 4 explanatory variables is

$$\hat{\sigma}^2 = \frac{\text{RSS}}{n - p} = \frac{132}{29 - 5} = 5.5$$

We calculate Mallows’  $C_p$  for each model.

$$C_p(0) = \frac{162}{29} = 5.586$$

$$C_p(1) = \frac{145 + 2(1)(5.5)}{29} = 5.379$$

$$C_p(2) = \frac{140 + 2(2)(5.5)}{29} = 5.586$$

$$C_p(3) = \frac{136 + 2(3)(5.5)}{29} = 5.828$$

$$C_p(4) = \frac{132 + 2(4)(5.5)}{29} = 6.069$$

The model with **1 explanatory variable** has the lowest  $C_p$  and is therefore the best.

- [8/8/2018] On page 813, in the solution to exercise 54.14, on the fourth line, delete the first  $\frac{15}{12}$ , the one without parentheses.
- [7/31/2018] On page 818, in the first displayed formula in Subsection 55.2.1, change  $\bar{X}$  to  $\bar{X}_j$ . In the second displayed formula, change  $\bar{x}_1$  to  $\bar{x}_j$  and put a right parenthesis after  $\bar{x}_j$ .
- [8/8/2018] On page 819, on the second line of Section 55.3, change “test data” to “training data”.
- [8/8/2018] On page 824, in the solution to exercise 55.11, on the second line, but a negative sign before  $\frac{1}{2} \frac{\partial f}{\partial \beta_0}$ .
- [10/28/2018] On page 825, in the solution to exercise 55.14, item 2, change “latter takes the response into account” to “former takes the response into account”.
- [10/25/2018] On page 840, in the box for exercises 57.2 and 57.3, in the table for centered moving averages, change the entry for 2011Q3 from 127.65 to 127.625.
- [7/24/2018] On page 852, in the solution to exercise 54.13, change the final answer to  $-1.25$ .
- [7/26/2018] On page 873, in the solution to exercise 61.1, change the final answer to 5.83614.
- [7/26/2018] On page 875, in the solution to exercise 61.14, on the first line, change 0.7 to 70 and  $+ 0.3$  to  $- 30$ .
- [10/18/2018] On page 891, change formula (63.7) to

$$\text{Forecast error} = s_w \sqrt{\sum_{i=0}^{t-1} a^{2i}}$$

- [10/18/2018] On page 892, in formula (63.7), change  $a^i$  to  $a^{2i}$ .
- [10/28/2018] On page 905, in the solution to exercise 64.5, on the second to last line, delete the “is derived” that is before “by”.
- [10/12/2018] On page 968, in question 6, in the two bullets, change “payment per loss” to “payment per payment”. Change the last line of the question to
- Calculate the percentage increase in the variance of payment per loss in 2016 over 2015.
- [7/26/2018] On page 991, in question 27, on the second line, change  $\beta_2$  to  $x_2$ .
- [7/26/2018] On page 991, in question 29, on the first bulleted line, insert a comma between 0.3111 and 0.5584.
- [10/28/2018] On page 1008, in the solution to question 34, on the first displayed line, change the denominator  $2\pi x^2$  to  $2\pi x^3$ .
- [10/28/2018] On page 1012, in the solution to question 11, on the last line, change  $\frac{l_{52}}{l_{51}}$  to  $\frac{l_{53}}{l_{52}}$ .
- [9/30/2018] On page 1029, replace the solution to question 41 with

The variance of the terms in the series, by formula (62.3), is

$$\sigma^2 = 4(1 + 0.8^2 + 0.6^2) = 8$$

The nonzero autocorrelations for the MA(2) series are at lags 1 and 2, based on formula (62.5).

$$\rho_1 = \frac{0.8 + (0.8)(0.6)}{1 + 0.8^2 + 0.6^2} = 0.64$$

$$\rho_2 = \frac{0.6}{1 + 0.8^2 + 0.6^2} = 0.3$$

Using formula (61.1), the variance of the sample mean is

$$\begin{aligned}\text{Var}(\bar{x}_t) &= \frac{\sigma^2}{n} (1 + 2(1 - 1/n)(\rho_1) + 2(1 - 2/n)(\rho_2)) \\ &= \frac{8}{10} (1 + 2(0.9)(0.64) + 2(0.8)(0.3)) = \boxed{2.1056} \quad (\text{E})\end{aligned}$$

[7/26/2018] On page 1038, in the solution to question 30, on the displayed line, change 1.4 in the exponent to 0.4.

[10/28/2018] On page 1056, in the solution to question 11, on the last line, 13.883 should be multiplied by 50,000 to obtain a final answer of  $\boxed{694,150}$ .

[10/28/2018] On page 1063, in the solution to question 45, on the second and fourth lines (once apiece), change  $\text{Cov}(x_t, x_{t+1})$  to  $\text{Cov}(x_t, x_{t+2})$ .

[8/12/2018] On page 1137, in the solution to question 33, on the second line, put “ln” before  $\frac{\mu}{1 - \mu}$ .

[7/5/2018] On page 1163, in the solution to question 38, on the first line, change “gamma” to “Gaussian”. On the fourth line, change “ $a(x) = \theta, b(\theta) = -(x - 1)^2/2x$ ” to “ $a(x) = -(x - 1)^2/2x, b(\theta) = \theta$ ”.

[9/13/2018] On page 1176, replace the solution to question 19 with

Since  $n/(n + 4) \rightarrow 1$  as  $n \rightarrow \infty$ , the limit of  $\mathbf{E}[\hat{\alpha}_n]$  is 7, making statement I true.

The mean square error of  $\hat{\alpha}_8$  is the square of the bias plus the variance:

$$\begin{aligned}\text{bias}_{\hat{\alpha}_8}(\alpha) &= \frac{7(8)}{8 + 4} - 7 = 2\frac{1}{3} \\ \text{Var}(\hat{\alpha}_8) &= \frac{140(7)}{15} = 65\frac{1}{3} \\ \text{MSE}(\hat{\alpha}_8) &= (2\frac{1}{3})^2 + 65\frac{1}{3} = 70\frac{7}{9}\end{aligned}$$

The variance approaches 70 from below as  $n \rightarrow \infty$ , since

$$\lim_{n \rightarrow \infty} \frac{n - 1}{2n - 1} = \frac{1}{2}$$

and

$$\frac{n - 1}{2n - 1} < \frac{n - 1}{2n - 2} = \frac{1}{2}$$

Since  $70\frac{7}{9} > 70$ , statement II is true.

Since the variance does not go to 0, we cannot be sure that  $\hat{\alpha}_n$  is consistent, so statement III is false.

**(B)**

[8/8/2018] On page 1182, in the solution to question 13, on the third line, change  $N \geq 4$  to  $N \geq 3$ . On the fifth line, change “ $N = 4$ . **(D)**” to “ $N = 3$ . **(C)**”. Delete the last sentence of the solution.

[8/8/2018] On page 1185, in the solution to question 32, on the third line, change “model 4” to “model 2”.